

Video Article

Coincubation Assay for Quantifying Competitive Interactions between *Vibrio fischeri* Isolates

Lauren Speare¹, Alecia N. Septer¹¹Department of Marine Sciences, University of North Carolina, Chapel HillCorrespondence to: Alecia N. Septer at asepter@email.unc.eduURL: <https://www.jove.com/video/59759>DOI: [doi:10.3791/59759](https://doi.org/10.3791/59759)Keywords: Immunology and Infection, Issue 149, interference competition, fluorescence microscopy, *Aliivibrio fischeri*, interstrain interactions, interspecies interactions, interbacterial killing

Date Published: 7/22/2019

Citation: Speare, L., Septer, A.N. Coincubation Assay for Quantifying Competitive Interactions between *Vibrio fischeri* Isolates. *J. Vis. Exp.* (149), e59759, doi:10.3791/59759 (2019).

Abstract

This manuscript describes a culture-based, coincubation assay for detecting and characterizing competitive interactions between two bacterial populations. This method employs stable plasmids that allow each population to be differentially tagged with distinct antibiotic resistance capabilities and fluorescent proteins for selection and visual discrimination of each population, respectively. Here, we describe the preparation and coincubation of competing *Vibrio fischeri* strains, fluorescence microscopy imaging, and quantitative data analysis. This approach is simple, yields quick results, and can be used to determine whether one population kills or inhibits the growth of another population, and whether competition is mediated through a diffusible molecule or requires direct cell-cell contact. Because each bacterial population expresses a different fluorescent protein, the assay permits the spatial discrimination of competing populations within a mixed colony. Although the described methods are performed with the symbiotic bacterium *V. fischeri* using conditions optimized for this species, the protocol can be adapted for most culturable bacterial isolates.

Video Link

The video component of this article can be found at <https://www.jove.com/video/59759/>

Introduction

This manuscript outlines a culture-based method to determine whether two bacterial isolates are capable of competitive interactions. When studying mixed populations, it is important to assess the extent to which the bacterial isolates interact, particularly whether isolates are directly competing through interference mechanisms. Interference competition refers to interactions where one population directly inhibits the growth or kills a competitor population¹. These interactions are important to identify because they can have profound effects on a microbial community's structure and function^{2,3}.

Mechanisms for microbial competition have been discovered broadly in genomes of bacteria from diverse environments including both host-associated and free-living bacteria^{4,5,6,7,8,9}. A variety of competition strategies have been described^{10,11} including diffusible mechanisms, such as bactericidal chemicals^{1,12} and secreted antimicrobial peptides¹³, as well as contact-dependent mechanisms that require cell-cell contact to transfer an inhibitory effector into target cells^{9,14,15,16,17,18}.

Although culture-based coincubations are commonly used in microbiology^{5,8,19}, this manuscript outlines how to use the assay to characterize the mechanism of competition, as well as suggestions for adapting the protocol for use with other bacterial species. Furthermore, this method describes multiple approaches for analyzing and presenting the data to answer different questions about the nature of the competitive interactions. Although the techniques described here were used previously to identify the interbacterial killing mechanism underlying intraspecific competition between symbiotic strains of coisolated *Vibrio fischeri* bacteria¹⁹, they are suitable for many bacterial species including environmental isolates and human pathogens, and can be utilized to evaluate both contact-dependent and diffusible competitive mechanisms. Steps in the protocol may require optimization for other bacterial species. Given that more model systems are expanding their studies beyond the use of isogenic organisms to include different genotypes^{10,16,20,21}, this method will be a valuable resource for researchers seeking to understand how competition impacts multi-strain or multi-species systems.

Protocol

1. Prepare Strains for Coincubation

1. Choose an appropriate reference strain that will serve as the target for bacterial competition during the coincubation assay. See **Discussion** for best practices when selecting a reference strain and how the reference strain will impact results. In this protocol, *V. fischeri* strain ES114 will serve as the reference strain.

2. Determining which selection and screening methods will be used to distinguish between the isolates in cocultivation

- Typically, transform strains with stable plasmids containing different antibiotic resistance genes (e.g. kanamycin or chloramphenicol) to select for each strain, as well as genes encoding different fluorescent proteins (e.g. GFP or RFP) for visually distinguishing strain types in coculture.
NOTE: Use of stable plasmids is required because if a strain loses the plasmid in the absence of selection, then this strain's numbers will be underestimated when quantified, and will not be visually detectable using fluorescence microscopy.
 - Tag cocultivated *V. fischeri* strains differentially such that one strain contains the plasmid pVSV102, which expresses a green fluorescent protein (GFP) and resistance to the antibiotic kanamycin (Kan^R), and the second strain contains plasmid pVSV208, which expresses a red fluorescent protein (dsRed) and resistance to the antibiotic chloramphenicol (Cm^R)²².
NOTE: Other differential selections can be used to distinguish cocultivating strains. See **Discussion** for additional methods.
 - Include the following control during initial optimization of the cocultivation assay to ensure the selection is robust. Plate each strain that is differentially tagged on agar plates containing the antibiotic that should select against it (i.e., plate strain 1 on media that selects for strain 2, and vice versa).
- Two days prior to cocultivation assay, streak reference strain pVSV102, reference strain pVSV208, and each competitor strain pVSV208 from -80 °C stocks onto LBS agar plates containing the appropriate antibiotic (e.g., 100 µg/mL kanamycin or 2 µg/mL chloramphenicol) and incubate overnight at 24 °C. Antibiotic selection is required to ensure all cells contain the plasmid at the beginning of the experiment.
 - One day prior to assay, pick and restreak four individual colonies per strain type (biological replicates) onto fresh LBS agar plates with appropriate antibiotic and incubated overnight at 24 °C.
NOTE: This step may require optimization for other bacterial species, as longer or shorter incubation times may be required to obtain sufficient cells depending on the growth rate of the organism of interest.

2. Cocultivate Bacterial Strains

1. Preparation of each bacterial strain for incubation (Figure 1A)

- Using a sterile toothpick, scrape the cells from the agar plate (as much as the size of a grain of rice) and re-suspend the cells in a 1.5 mL microcentrifuge tube containing 1 mL of LBS broth. Break up the cell clumps by pressing them into the side of the tube and pipetting up and down vigorously.
 - Cap the tube and vortex for 1-2 s. If the cell clumps are still visible, continue to vortex or pipette up and down until the sample is visually uniform.
 - Repeat steps 2.1.1-2.1.2 for all samples.
- Measure and record the optical density at 600 nm (OD₆₀₀) for all samples. Samples will likely need to be diluted up to ten-fold using LBS broth to obtain an accurate OD measurement. Normalize each sample to desired OD₆₀₀. Strains are typically normalized to an OD₆₀₀ ~1.0, which translates to ~10⁸ bacteria/mL for *V. fischeri*.
NOTE: This step may require optimization for different bacterial species as the inoculum cell density impacts cocultivation results depending on the mechanism of competition. See discussion for optimization.
- Cocultivating strains (Figure 1B, C)**
 - Mix the reference strain and competitor strain in a 1:1 ratio (v/v) by adding 100 µL of each strain (normalized to OD 1.0) to a labeled 1.5 mL centrifuge tube. As a control, mix the reference strain with a differentially tagged version of itself (reference strain pVSV102 + reference strain pVSV208). This control is required for later statistical analysis to determine whether the presence of a competitor strain impacts the growth of the reference strain. Vortex the mixed-strain culture for 1-2 s.
NOTE: This step may require optimization as the starting ratio can significantly impact results and may need to be adjusted. See discussion for optimization.
 - Repeat step 2.3.1 for each biological replicate. After completing this step, there should be a total of eight mixed-strain tubes: four biological replicates with differentially-tagged reference strains (control), and four biological replicates with differentially-tagged reference and competitor strains (experimental).
 - Spot 10 µL of each control and experimental mixture onto Petri plates containing LBS agar; these culture spots will be used for fluorescence microscopy after the incubation.
 - Allow the spots to dry completely on the bench until all liquid has been absorbed into the agar and incubate the Petri plates at 24 °C for 24 h. A minimum of 15 h is required for *V. fischeri* strains tagged with pVSV102 and pVSV208 to grow to a high enough cell density to visualize GFP and RFP, respectively, at the population level using a fluorescence dissecting microscope. Here, we use a 24 h incubation for imaging mixed spots.
NOTE: It is important to use plates that are not too moist or too dry. If the plates are too moist, cocultivation spots will not be absorbed into the agar plate; avoid using plates poured on the same day. If the plates are too dry, small waves or cracks may form on the agar surface, making cocultivation spots irregular in shape.
 - Using the same bacterial suspensions as in step 2.3.3, spot 10 µL of each control and experimental mixtures into 24-well plates containing 1 mL of LBS agar per well. As in step 2.3.3, ensure the agar in 24-well plates has the appropriate moisture. Allow the spots to dry completely and incubate at 24 °C for 5 h. These culture spots will be used to quantify the colony forming units (CFUs) for each strain at the end of the experiment.
NOTE: Spotting bacterial suspensions for growth in individual wells allows for easier resuspension at the end time point. This step may be accomplished using square sterile filter pieces as a more economical approach to using 24-well plates. Square sterile filter pieces can be placed on an agar plate and 10 µL of each experimental mixture can be spotted onto these filters, rather than onto 24-well plates. After the cocultivation time, the filters can be transferred into a 1.5 mL centrifuge tube and the cocultivation spot can be resuspended by vortexing or pipetting up and down. A 5 h incubation time is sufficient to detect interbacterial killing between *V. fischeri* isolates¹⁹, however, this cocultivation time may need to be shorter or longer for other species or competitive mechanisms. See discussion for more details.

4. Serial dilution for starting inoculum

- To perform a serial dilution of starting coinoculation mixtures, rotate a 96-well plate 90° so there are eight columns and twelve rows. Each row will contain a sample of the undiluted mixture in the first column followed by seven ten-fold serial dilutions across the remaining wells in the row (**Figure 2A**).
- Label each row with the strain ID, the replicate number, and the time (starting inoculum = T₀). Label the LBS agar plates supplemented with the appropriate antibiotic on which to spot the dilution series. Be sure to identify which strain each antibiotic plate is selecting for.
- Using a multichannel pipette, add 180 µL of LBS broth to each well, leaving the first column empty for the undiluted coinoculation mixture.
NOTE: Researchers may choose to use phosphate buffered saline (PBS) to resuspend coinoculation spots and perform serial dilution if working with a particularly fast growing bacterial species or consistently performing large experiments where serial dilutions may take a long time to perform. Using PBS in these instances will prevent significant outgrowth of bacteria during the process of performing serial dilutions. However, it is important to be consistent with which solution is used (e.g., PBS or LBS) across all experiments, regardless of their size or duration.
- Using a 200 µL single channel pipette, transfer 100 µL of the coinoculation mixture from the tube to the first column well. Discard the tip and repeat for all coinoculation mixtures.
- Using a multichannel pipette, transfer 20 µL from column 1 to 2 and mix by pipetting up and down. Discard the tips and repeat for each column so that by the end, each row contains a ten-fold serial dilution of the initial coinoculation mixture.
NOTE: Be consistent with the number of times dilutions are pipetted up and down. For example, depress the pipette handle five times for each dilution to be consistent across the dilution series.
- Using a multichannel pipette fitted with eight tips, aspirate 5 µL from each well in a dilution series (i.e., one row of eight wells) and spot it onto the LBS agar plate that selects for the reference strain (supplemented with kanamycin). Repeat this step spotting onto the plate selecting for the competitor strain (supplemented with chloramphenicol) (**Figure 2B**). Allow the spots to dry completely prior to placing in incubator.
NOTE: The same tips may be used to spot a replicate dilution series on plates selecting for the reference and competitor strain, but must be changed between replicates. Avoid touching the end of the pipette tip to the LBS agar plates as this can create a depression that can resemble a bacterial colony and skew results during plate counts. If the tip does touch the plate, make a note and do not include the depression in colony counts.

5. Taking T final measurements

- After the coinoculation spots in the 24-well plates have been incubated for 5 h at 24 °C, add 1 mL of LBS broth to each well and resuspend the coinoculation spots by pipetting up and down until all cells are resuspended. Be consistent with the vigor in which the cells are resuspended across all replicates. For example, depress the pipette handle eight times to fully resuspend each sample.
- Once each coinoculation spot is resuspended, prepare a 96-well plate for a serial dilution and LBS agar plates with the appropriate antibiotics on which to spot the dilution in the same way as step 2.4.1.
- Perform steps 2.4.2-2.4.6 to complete the serial dilution for T final measurements.
NOTE: An important control should be included during initial optimization of the coinoculation assay. At the end of the coinoculation period, plate the mixed strains onto media that includes both antibiotics. Neither strain should be able to grow unless 1) a spontaneous mutation occurs, or 2) selectable markers are exchanged between strains.

3. Visualizing Coinoculations Using Fluorescence Microscopy

- Image each coinoculation spot on LBS agar petri plates from steps 2.3.3 and 2.3.4 using a stereo microscope equipped for green and red fluorescence detection, which correspond with the fluorescence proteins being expressed on the stable plasmids.
- Begin by taking images of the control coinoculation spot (reference strain pVSV102 + reference strain pVSV208).**
 - Adjust magnification so the spot is in focus.
 - Using the appropriate excitation light and filter to observe GFP, adjust the exposure time so only fluorescence from the coinoculation spot is detectable and any background fluorescence from the media is low or not visible.
 - Change the excitation light and filter to observe RFP, adjusting the exposure time to minimize background from the medium.
- For each coinoculation spot from the experimental treatments, image for the reference strain using the GFP filter and exposure time determined in step 3.2.2.
- Image the competitor strain using the RFP filter, adjusting the exposure time so that fluorescence is only observed from the coinoculation spot and background fluorescence from the medium is minimal. Save both images and name them to include both strain IDs, the replicate number, the incubation time when the image was taken (e.g., 24 h). Repeat for all replicates.
NOTE: Coinoculation time may need to be adjusted for different plasmids or bacterial species. See discussion for optimization.

4. Data Analysis

- Calculating CFUs for each control and experimental treatment at each timepoint (T start and T final)**
 - Once the colonies are visible on serial dilution plates (e.g., 24 h at 24 °C for *V. fischeri*), identify the dilution factor where individual colonies can be counted and have not grown together into large, multi-colony clusters (**Figure 2B**). For each replicate, count and record the number of individual colonies for a given dilution. This number represents the CFUs for that replicate.
 - Convert CFUs for the dilution to total CFUs for each strain in the coinoculation using the formula below:
[CFUs ÷ (dilution factor x vol. of dilution spot)] x vol. of the undiluted sample = total CFUs
Example: T₀: [(6 CFUs) ÷ (10⁻⁶ x 0.005 mL)] x [0.01 mL] = 1.2E7 total CFUs of strain 1 in the mixed spot at the beginning of the experiment
T₅: [(4 CFUs) ÷ (10⁻³ x 0.005 mL)] x [1.0 mL] = 8.0E5 total CFUs of strain 1 in the mixed spot after 5 h coinoculation

The CFU values for each strain at each time point are the raw data that can be used for several different analyses and statistical tests (see section 4.2 below).

2. **Perform the following data transformations to determine whether a competitor strain outcompetes the reference strain by: (i) determining whether the proportion of the reference strain decreases in the presence of the competitor strain, or (ii) calculating the log relative competitive index (RCI). These analyses convert raw data into ratios and can be useful to determine the competitive outcome of an interaction, but cannot determine the mechanism of competition (i.e., killing vs growth inhibition).**

1. **Calculating the proportion of each strain for each time point during the coinubation**

1. Convert CFU data to the proportion of each strain in a treatment. For the reference strain (RS) at T0, divide the total CFUs for the reference strain at T0 by the sum of total CFUs for the reference strain and the competitor (CS) at T0:

$$RS_{T0} \text{ CFUs} \div (RS_{T0} \text{ CFUs} + CS_{T0} \text{ CFUs}) = \text{Proportion of reference strain at T0.}$$
 Perform the same calculation to determine the proportion of the competitor strain at T0:

$$CS_{T0} \text{ CFUs} \div (RS_{T0} \text{ CFUs} + CS_{T0} \text{ CFUs}) = \text{Proportion of competitor strain at T0}$$
 Repeat this process for each time point and replicate for both the experimental treatment and the control treatment (differentially-tagged reference strain coinubation).
2. Graph the average proportion of each strain type at T0 and T5 in a stacked bar graph (**Figure 3A**).
3. Ensure that the desired starting ratio of strains was obtained. For coinubations where strains were initially mixed 1:1, each strain type should comprise ~0.5 of the population at T0 and should not be statistically different from each other using a Student's t-test ($P > 0.05$). If the proportion of one strain is significantly larger than that of the other strain at T0, then repeat the experiment until a 1:1 starting ratio is achieved.
4. Determine whether the competitor strain comprises a significantly larger proportion of the population after 5 h. Perform a Student's t-test comparing the proportion of each strain in the experimental treatment at T0 and T5. If the proportion of competitor strain is significantly different from that of the reference strain at T5 ($P < 0.05$), this result suggests strain competition may have occurred. Proceed to step 4.2.1.5.
5. To determine whether the presence of the competitor strain reduced the proportion of the reference strain after 5 h, perform a Student's t-test comparing the proportion of the reference strain in the control with the proportion of the reference strain in the experimental treatment (reference strain v competitor strain).

NOTE: If the proportion of the reference strain is statistically lower in the experimental treatment relative to the control ($P < 0.05$), then the competitor strain significantly reduced the proportion of the reference strain after 5 h and outcompeted the reference strain.

2. **Calculating the log RCI value for each treatment (including the control)**

NOTE: The relative competitive index, or RCI, is a single value that compares how the ending ratio of strain types differs from the starting ratio. The RCI value is log-transformed such that a log RCI value greater than zero indicates the competitor strain (CS) outcompeted the reference strain (RS), a log RCI value less than zero indicates the reference strain outcompeted the competitor strain, and a log RCI value of zero indicates that neither strain was outcompeted.

1. Calculate log RCI values for each control and experimental treatment using the following equation:

$$\text{Log} [(CS_{T5} \text{ CFU} \div RS_{T5} \text{ CFUs}) \div (CS_{T0} \text{ CFU} \div RS_{T0} \text{ CFUs})] = \text{log RCI}$$

2. Graph log RCI values using a separated box & whiskers plot where data are graphed in the horizontal direction (**Figure 3B**).
3. Determine whether each treatment was significantly different from zero by performing a student's t-test comparing the log RCI values of each treatment (including the control) to a data set comparing the same number of replicates all with a zero value. If the log RCI values for the experimental treatment are statistically greater than zero ($P < 0.05$), then the competitor strain outcompeted the reference strain.

NOTE: The control should not be statistically different from zero ($P > 0.05$), because the differentially tagged reference strains are isogenic.

4. Perform a Student's t-test to determine whether log RCI values for the experimental treatment were significantly greater than the control ($P < 0.05$). If the log RCI values from the experimental treatment are statistically greater than the control treatment, then the competitor strain outcompeted the reference strain.

3. **Perform the following statistical analysis to determine the mechanism by which the competitor strain outcompetes the reference strain: (i) analyze raw total CFU data, or (ii) examine percent recovery of the reference strain. These analyses can be more cumbersome to view relative to those from step 4.2, but will identify whether the competitor strain outcompetes the reference strain by outgrowing it, inhibiting reference strain growth, or actively eliminating the reference strain.**

1. **Analyzing raw total CFU data**

1. Graph raw total CFU data for both strains using an interleaved scatter plot, where replicates are shown as individual data points (**Figure 4A**). Display data on a log scale, and add a horizontal line indicating the average T0 CFUs for both strains.
2. To determine whether the presence of the competitor strain negatively affected the reference strain, perform a Student's t-test comparing reference strain CFUs for the control and experimental treatments at T5. If the reference strain CFUs in the experimental treatment are statistically lower than in the control ($P < 0.05$), then the presence of the competitor strain significantly affected the reference strain.
3. To determine whether the presence of the competitor strain either inhibited the growth of or eliminated the reference strain, perform a Student's t-test comparing the reference strain CFUs at T0 and T5 for the control and experimental treatments.
NOTE: In the absence of a competitor, the reference strain should show a significant increase in CFU when comparing T0 and T5 for the control treatment. When analyzing the experimental treatment, if reference strain CFUs at T5 are not statistically different compared to T0 ($P > 0.05$), then the competitor strain inhibited the growth of the reference strain. If reference strain T5 CFUs are significantly lower than T0 CFUs ($P < 0.05$), then the competitor strain killed the reference strain.

2. **Calculating percent recovery of the reference strain**

1. Transform total CFU data for the reference strain (RS) into percent recovery data using the below equation:

2. $(RS_{T5} \text{ CFUs} + RS_{T0} \text{ CFUs}) \times 100 = \% \text{ recovery of reference strain}$
3. Display the percent recovery of reference strain using a separated bar graph where each bar represents either the control or experimental treatment (**Figure 4B**). Add a dashed line at $y = 100$ to indicate 100% recovery of reference strain (no increase or decrease in reference strain CFUs from 0 h to 5 h).
4. To determine whether the competitor strain negatively affected the reference strain, perform a Student's t-test comparing reference strain percent recovery for the control treatment to the experimental treatment. If the percent recovery is significantly less than the control ($P < 0.05$), then the competitor strain negatively affected the reference strain.
5. To identify whether the competitor strain inhibited the growth of or eliminated the reference strain, perform a Student's t-test comparing reference strain percent recovery for the experimental treatment and a data set with the same number of replicates all with a value of 100.
NOTE: If the experimental treatment is not statistically different from 100 ($P > 0.05$), then the competitor strain inhibited the growth of the reference strain. If the reference strain percent recovery in the experimental treatment is significantly lower than 100 ($P < 0.05$), then the competitor strain killed the reference strain.

3. Interpreting fluorescence microscopy images

1. Once fluorescence microscopy images are taken, determine whether each strain is visibly detectable in the coincubation spot. For the control coincubation (reference strain pVSV102 + reference strain pVSV208), both GFP and RFP should be visible from one coincubation spot. If this is not the case, set up the experiment again ensuring strains are properly labeled and coincubated on antibiotic-free plates.
2. For each experimental coincubation spot, check for possible outcomes.

NOTE: If the competitor strain is visible, but the reference strain is not detected, that suggests the competitor strain killed or inhibited the growth of the reference strain. If both strains are present and uniformly mixed (GFP and RFP present throughout the coincubation spot), these data suggest the competitor strain does not compete with the reference strain and both strains coexist. If both strains are present but are spatially separated (microcolonies of either GFP or RFP are present throughout the coincubation spot), that suggests the reference strain and competitor strain may be engaging in competitive interactions and modifications to the reference strain or initial coincubation ratio may be required. See discussion for more details.

5. Determining Whether Interaction is Contact-dependent

NOTE: If you find that one strain kills or inhibits the reference strain, the interaction may be diffusible or contact-dependent. To determine whether the interaction is dependent on cell-cell contact, perform a coincubation assay as described above for steps 1-2 with the following modifications.

1. Perform steps 1.1-2.2.
2. Once strains are normalized, physically separate strains using a nitrocellulose filter with a 0.22 μm pore size. This pore size allows for the diffusion of antimicrobials and small molecules but prevents physical contact between *V. fischeri* cells.
NOTE: If the interaction is dependent on cell-cell contact, separation of the reference strain from the competitor strain with the filter should abrogate the killing or inhibitory phenotype and the reference strain should not have reduced CFUs. If the interaction is not dependent on cell-cell contact, and is a diffusible mechanism, separating the strains should not prevent the competitor strain from killing the reference strain.
 1. Spot 5 μL of the reference strain onto the center of a filter and spot 5 μL of the competitor strain onto the center of a different filter. Place the filter containing the reference strain onto the surface of an LBS agar plate. Place the filter containing the competitor strain directly on top of the filter with the reference strain.
NOTE: Each strain is spotted onto filters rather than the bottom strain spotted directly onto the agar plate so strains can be more easily placed directly on top of one another.
 2. If there is concern that strains are not stacked directly on top of one another, decrease the volume of reference strain inoculum spotted onto the filter. This will ensure the area of the reference strain spot is smaller and fully above or beneath the competitor strain. Alternate which strain is on top and on bottom to account for differences in diffusion of nutrients from the agar medium.
 3. To ensure the filter allows for diffusion of small molecules, include a control treatment where the reference strain is spotted onto a filter and an antibiotic that it is sensitive to (chloramphenicol) is spotted onto the other. Stack the filters directly on top of one another and place on an LBS agar plate. The antibiotic should be able to diffuse through the filter and should kill the reference strain.
3. Incubate the plates with filters at 24 $^{\circ}\text{C}$ for 5 h. Do not invert the agar plate when incubating to avoid moving the filters.
4. Perform serial dilutions for the starting inoculum according to step 2.4.
5. **Taking T final measurements**
 1. Using sterile tweezers, transfer each set of filters to a 50 mL Falcon tube containing 5 mL of LBS broth. Ensure that the filters are physically separated within the tube to allow for full resuspension of each strain type.
 2. Resuspend the strains from filters by pipetting up and down until the filters are clear of all cells. Use tweezers to rotate the filters and clear cell material from both sides. Sterilize tweezers with ethanol between samples.
 3. Perform serial dilution for T final according to step 2.5. When calculating total CFUs for T final, adjust "total volume of undiluted sample" from 1 mL to 5 mL.

Representative Results

In order to assess competitive interactions between bacterial populations, a coinoculation assay protocol was developed and optimized for *V. fischeri*. This method utilizes stable plasmids that encode antibiotic resistance genes and fluorescent proteins, allowing for differential selection and visual discrimination of each strain. By analyzing the data collected from the coinoculation assay, the competitive outcome of an interaction and the mechanism of the interaction can be identified. As an example, the following experiments were performed using *V. fischeri* isolates. Coinoculated strains harbored one of two stable plasmids: pVSV102 expressing kanamycin resistance and GFP+, or pVSV208 expressing chloramphenicol resistance and DsRed+. In the sample data, strains were mixed in a 1:1 ratio and incubated on LBS agar plates for 5 h. As a control treatment, differentially tagged versions of the reference strain were coinoculated with each other. The experimental treatments were performed with the reference strain (harboring pVSV102) and either competitor strain 1 or competitor strain 2 (harboring pVSV208). Cultures of each strain were prepared and coinoculated as described above and as shown in **Figure 1** and **Figure 2**.

In **Figure 3**, data analyzed to determine whether competitor strain 1 or 2 outcompeted the reference strain are presented in two ways. In **Figure 3A**, the proportion of each strain type for each time point during the experiment was calculated according to step 4.2.1. In the experimental treatments, the sample data show the reference strain and competitor strain 1 are present at a proportion of 0.5 at the beginning (0 h) and end (5 h) of the experiment, which is consistent with what is observed in the control treatment. These data show the proportion of the strains did not change after a 5 h coinoculation, and therefore no competition was observed. By contrast, when the reference strain was incubated with competitor strain 2, the reference strain was present at a proportion of 0.5 at the beginning (0 h), and a proportion <0.01 at the end (5 h) of the experiment, which was significantly lower than the control treatment (Student's t-test: $P < 0.001$). These data indicate that the proportion of the reference strain decreased in the presence of competitor strain 2, and therefore suggests competition between competitor strain 2 and the reference strain occurred. This type of analysis should be applied when determining how the proportion of strains within a community changes over time but cannot be used to determine the mechanism of the competitive interaction, and therefore should be combined with additional analysis. For example, the proportion of the reference strain decreasing in the presence of competitor strain 2 could be attributed to several types of interactions: (i) strain 2 grew more quickly than the reference strain, (ii) strain 2 inhibited the growth of the reference strain, or (iii) strain 2 eliminated the reference strain through killing.

In **Figure 3B**, the log relative competitive index (RCI) was calculated for each treatment according to step 4.2.2. When the reference strain was incubated with competitor strain 1, log RCI values were not statistically different from zero or from the control treatment ($P > 0.05$), suggesting competition between strains was not observed. When the reference strain was incubated with competitor strain 2, however, log RCI values were significantly greater than zero and the control treatment (Student's t-test: $P < 0.001$). These data suggest strain 2 outcompeted the reference strain. Analyzing log RCI values provides a simple method to determine whether one strain outcompeted the other during the incubation period. Because this analysis incorporates the ratio of strains at the end (5 h) and the beginning (0 h) of the experiment, the starting ratio can dramatically impact the result. Therefore, the starting ratio should be examined and considered when deriving conclusions from log RCI data. Furthermore, this analysis does not provide information about the competitive mechanism and simply reports how the ratio of strains change during the incubation.

Figure 4 displays two methods of data analysis to determine the mechanism of competition for a given interaction. In **Figure 4A**, total CFUs for each strain at each time point of the experiment are displayed. When the reference strain was incubated with competitor strain 1, CFUs of both strains increase over the course of 5 h and CFUs for the reference strain were not significantly different from strain 1 or the control at 5 h ($P > 0.05$). These data indicate that the reference strain grew in the presence of strain 1, and suggest no competition occurred. However, when the reference strain was incubated with competitor strain 2, strain 2 CFUs increased after 5 h but CFUs for the reference strain decreased. Reference strain CFUs were significantly lower than strain 2 CFUs and the control at 5 h (Student's t-test: $P < 0.002$). These data indicate that the reference strain CFUs decrease in the presence of strain 2, and suggest strain 2 kills reference strain cells. If the reference strain did not show a decrease in CFUs, but rather no change (no statistical difference between reference strain CFUs at 0 h and 5 h), these data would suggest strain 2 outcompeted the reference strain by inhibiting the growth of the reference strain. Analyzing untransformed total CFU data is particularly informative, as CFUs for both strains at each time point are displayed independently and can be used to identify the mechanism of competition.

Figure 4B shows the percent recovery of the reference strain in order to determine how the presence of a competitor strain affects the reference strain. When the reference strain was incubated with competitor strain 1, a ~3,200 percent recovery was observed, which was not statistically different from the control and indicates strain 1 did not affect the percent recovery of the reference strain. When the reference strain was incubated with competitor strain 2, a ~4 percent recovery was observed, which was significantly lower than the control (Student's t-test: $P < 0.002$). The percent recovery was also significantly less than 100 (Student's t-test: $P < 0.002$), indicating strain 2 outcompeted the reference strain by killing reference strain cells. If the percent recovery was not statistically different from 100, those data would suggest strain 2 inhibited the growth of the reference strain. Percent recovery data provides a simplified way to characterize the mechanism of competition by examining how the reference strain population responds to the presence of a competitor strain. However, displaying the data in this way excludes information about the starting ratio as well as how the abundance of the competitor strain changed throughout the incubation.




Figure 1: Flowchart illustrating the coincubation assay. (A) Bacterial strains harboring either pVSV102 (reference strain indicated Ref. Strain or R.S.) or pVSV208 (competitor strain indicated Comp. Strain or C.S.) are grown separately on media selective for either the reference strain (LBS Kan) or competitor strain (LBS Cam). Strains are then resuspended in LBS broth and normalized to an OD = 1.0. (B) The reference strain and competitor strain are mixed at a 1:1 ratio by volume. A serial dilution is performed with this mixture to determine CFUs for both strains at 0 h. (C) The strain mixture is then spotted onto 24-well plates containing LBS agar. Each replicate is spotted into its own well. Spots are allowed to dry and then incubated at 24 °C for 5 h. After 5 h, a serial dilution is performed to quantify CFUs for each strain. (D) The strain mixture from panel B is also spotted onto LBS agar Petri plates allowed to dry and incubated at 24 °C for 24 h. At 24 h, the coincubation spot is imaged using a fluorescence dissecting microscope that is adapted to detect green (reference strain) and red (competitor strain) fluorescence. Scale bar = 1 mm. [Please click here to view a larger version of this figure.](#)




Figure 2: Representative images of plates required for a serial dilution. (A) 96-well plate used to perform serial dilution. The plate is rotated such that there are 12 rows and 8 columns. Descriptors of each treatment include the strains used and the plasmids they harbor (e.g., Reference strain with pVSV102 and Competitor strain with pVSV208) and the replicate number for each row (R1, R2, R3, or R4). The first column is the undiluted sample and each column to the right represents a 10-fold dilution from the previous column (dilution factor listed above). (B) LBS agar plate used to determine CFUs for the reference strain on LBS Kan plates (top) and competitor strain on LBS Cam plates (bottom) from the experimental treatment. Each row is one replicate in a treatment (e.g., R1) and the dilution factor of each spot is listed at the top of the plate. The number of CFUs counted for each replicate is listed to the right. [Please click here to view a larger version of this figure.](#)




Figure 3: Sample data for assessing whether competitor strains outcompete the reference strain. (A) The proportion of co-culturing strains harboring either pVSV102 (dark gray) or pVSV208 (light gray). R.S. indicates the reference strain and C.S. indicates the competitor strain. Dashed horizontal line indicates a proportion of 0.5. Asterisk indicates the reference strain made up a statistically smaller proportion of the population than the competitor strain or reference strain in the control at 5 h (Student's t-test: $P < 0.001$); ns indicates not significant ($P > 0.05$). **(B)** Log relative competitive index (RCI) for co-culturing assays. Dashed vertical line indicates $\log RCI = 0$. Asterisk indicates the $\log RCI$ value is statistically greater than zero and the control (Student's t-test: $P < 0.001$). Error bars indicate SEM. [Please click here to view a larger version of this figure.](#)




Figure 4: Sample data for determining the mechanism of competition. (A) Total CFU counts for coinubation assays performed with *V. fischeri* isolates that were differentially tagged with either pVSV102 or pVSV208. CFUs were collected at the beginning of the experiment (0 h) and after 5 h incubation. R.S. indicates reference strain and C.S. indicates competitor strain. Dashed horizontal line indicates the average 0 h CFUs for both strains; asterisk indicates reference strain CFUs were statistically lower in the experimental treatment relative to the control treatment at 5 h (Student's t-test: $P < 0.002$). (B) Percent recovery of the reference strain. Horizontal dashed line indicates 100% recovery (no increase or decrease in CFUs); asterisk indicates percent recovery was statistically lower than 100% and the control treatment (Student's t-test: $P < 0.002$). Error bars indicate SEM. [Please click here to view a larger version of this figure.](#)




Figure 5: Sample data for incubation time and imaging optimization. (A) Total CFUs for coinubation assays where CFUs were collected at the beginning of the experiment (0 h), after 5, 12, and 24 h incubation. R.S. indicates reference strain and C.S.2 indicates competitor strain 2. Dashed horizontal line indicates the average 0 h CFUs for both strains; asterisks indicate reference strain CFUs were statistically lower in the experimental treatment relative to the control treatment at the given time point (Student's t-test: $P < 0.002$). Error bars indicate SEM. (B) Fluorescent microscopy images corresponding with CFU data in panel A. Scale bar = 1 mm. [Please click here to view a larger version of this figure.](#)




Figure 6. Sample data for coinoculation ratio optimization. Coinoculation experiments were performed between the reference strain (R.S.) harboring pVSV102 (green, top row) and competitor strain (C.S.) harboring pVSV208 (red, bottom row) and fluorescent microscopy images were taken at 24 h. (A) Strains were mixed in a 1:1 ratio or (B) a 1:5 ratio where the reference strain, containing pVSV102, was outnumbered by the competitor strain containing pVSV208. Scale bar = 1 mm. [Please click here to view a larger version of this figure.](#)

Discussion

The coinoculation assay described above provides a powerful method to discover interbacterial competition. This approach allowed for the identification of intraspecific competition among *V. fischeri* isolates and characterization of the competitive mechanism¹⁹. Although the method described was optimized for the marine bacterium *V. fischeri*, it can be easily modified to accommodate other bacterial species including clinical and environmental isolates. It is important to note that competitive mechanisms are often conditionally regulated^{5,6,23,24,25,26,27,28}, thus small differences in growth conditions (e.g., shaking vs standing culture, temperature, etc.) and media type (e.g., salt content) can dramatically affect the results. Therefore, optimizing coinoculation conditions is likely necessary for different bacterial species as well as different competitive mechanisms. It is best to choose culture conditions that closely reflect the natural environment of the isolates. For example, coinoculation assays between *V. fischeri* strains were performed with LBS media at 24 °C, to reflect the salinity and temperature of the marine environment. However, some bacteria are naturally competent in their environment^{27,28} and therefore could take up genetic material released by lysed cells during antagonistic interactions²⁹. To prevent such DNA transfer from impacting coinoculation results, it is important to use conditions that do not promote competence or strains that are not competent, either naturally or through inactivation of DNA uptake machinery. Moreover, experimental parameters such as cell growth phase, culture density, incubation time, or starting strain ratio may also require optimization for different bacterial species or competitive mechanisms. For example, initial culture density will dictate the amount of cell-cell contact between strains, which can affect the ability of bacteria to deploy contact-dependent mechanisms of competition.

Figure 5A displays the process of optimization of coinoculation assays with *V. fischeri* isolates. Here, a range of incubation times for CFU collection and fluorescent microscopy imaging were evaluated to determine the optimal time for each metric to be collected. CFUs were collected immediately after the mixture of the reference strain and competitor strain 2 was spotted onto LBS agar plates (0 h), and CFU measurements and fluorescent microscopy images were taken immediately and after 5, 12, and 24 h. These sample data highlight the importance of thorough optimization prior to drawing any conclusions about the interaction between two strains. For example, two different conclusions about the mechanism of interaction can be deduced based on when CFUs are collected: CFUs from 5 or 12 h indicate strain 2 killed the reference strain, while CFUs collected at 24 h suggest strain 2 inhibits the growth of the reference strain.

The optimal time for visualization of coinoculation spots through fluorescent microscopy may be different than the optimal time for CFU collection. **Figure 5B** displays fluorescent microscopy images of coinoculation spots at 0, 5, 12, and 24 h. At 0 and 5 h, the coinoculation spots are not visible with fluorescent microscopy. For images taken at 12 h, both strains in the control treatment are visible, yet the RFP (reference strain harboring pVSV208) is notably dimmer. In the experimental treatment at 12 h, competitor strain 2 is visible (yet dim) and the reference strain is not detectable. Strain-specific differences between bacterial isolates can affect the expression of fluorescent proteins, and thus brightness of the cells in the mixed spot. Because RFP is notably dimmer than GFP in the control, the coinoculation spots should continue to be incubated and be

imaged again at a later time. In images taken at 24 h, both strains are visibly detectable and at a similar brightness in the control experiment. In the experimental treatment strain 2 is visible while the reference strain is not observed within the coinubation spot. 15 - 24 h incubation time is sufficient to visualize GFP and RFP for *V. fischeri* using stable plasmids pVSV102 and pVSV208, respectively, but the incubation time may need to be adjusted for different plasmids or bacterial species. Although the optimal time for visualization of coinubation spots and collecting CFU data are different, imaging at 24 h is a good way to quickly screen interactions for *V. fischeri*, because the result obtained from imaging at 24 h (target is visible or not) reflects the more time-intensive quantitative data obtained from plating CFUs at 5 or 12 h.

The starting ratio can significantly impact results, particularly when incubating two inhibitory strains, and may need to be adjusted to account for strain specific differences in killing efficiency or growth rate. For example, **Figure 6A** displays fluorescent microscopy images of experiments where the reference strain was coinubated with itself (control) and three other *V. fischeri* isolates starting at a 1:1 ratio. In these sample data, the reference strain is visibly detected when incubated with itself, competitor strain 1, and competitor strain 3 after 24 h. However, when the starting ratio was adjusted to 1:5 (i.e., 50 μ L of reference strain mixed with 250 μ L of competitor strain) the reference strain is only visibly detected when coinubated with itself and strain 1, indicating that both strain 2 and strain 3 outcompete the reference strain. This adjustment prevents the faster growth rate of the reference strain from obscuring the effect of any interference competition mechanisms exhibited by the competitor strains. Based on the results in **Figure 6A**, a ratio of 1:5 (reference strain : competitor strain) should be used to screen additional *V. fischeri* strains for the ability to kill the reference strain.

This protocol discriminates between coinubating strains by differentially labeling strains with plasmids containing either kanamycin or chloramphenicol resistance genes. However, different antibiotics or other selection methods may be better suited for different bacterial species. Other methods for differential selection could include: 1) exploiting a strain/species-specific auxotrophy for specific growth factors (e.g., DAP or thymidine), 2) conditional growth requirements (e.g., one strain grows at 37 °C while the other does not), or 3) counterselection markers that eliminate or inhibit the growth of the tagged strain when grown under appropriate conditions to express a "kill" gene (e.g., *ccdB* or *sacB*).

Selecting the appropriate reference strain is critical for obtaining and interpreting reproducible results from the coinubation assay. A reference strain should be well-studied (i.e., have a broad body of scientific literature), have no apparent killing or inhibitory ability, and ideally have a sequenced genome. For example, certain strains of *Escherichia coli* are common reference strains for many bacterial coinubation experiments^{30,31}. However, *E. coli* may not be ecologically relevant for a given competitive mechanism or competitor, which can affect results. For example, some bacteria may have evolved mechanisms specifically targeting closely-related species or competitors for the same ecological niche and their competitive mechanism would not be effective against an *E. coli* reference strain.

In summary, the method described here aims to provide an easily modified and robust approach to evaluate interbacterial interactions and competition. This method can be applied to bacterial isolates relevant to environmental or clinical research, and can be used to explore diverse mechanisms of microbial interaction that have been previously unknown or difficult to investigate.

Disclosures

The authors have nothing to disclose.

Acknowledgments

We would like to thank reviewers for their helpful feedback. A.N.S. was supported by the Gordon and Betty Moore Foundation through Grant GBMF 255.03 to the Life Sciences Research Foundation.

References

- Hibbing, M.E., Fuqua, C., Parsek, M.R., Peterson, S.B. Bacterial competition: surviving and thriving in the microbial jungle. *Nature Reviews Microbiology*. **8**(1), 15-25 (2010).
- Nyholm, S.V., McFall-Ngai, M. The winnowing: establishing the squid-Vibrio symbiosis. *Nature Reviews Microbiology*. **2**(8), 632-642 (2004).
- Dörr, N.C.D., Blockesh, M. Bacterial type VI secretion system facilitates niche domination. *Proceedings of the National Academy of Sciences of the United States of America*. **115**(36), 8855-8857 (2018).
- MacIntyre, D.L., Miyata, S.T., Kitaoka, M., Pukatzki, S. The Vibrio cholerae type VI secretion system displays antimicrobial properties. *Proceedings of the National Academy of Sciences of the United States of America*. **107**(45), 19520-19524 (2010).
- Salomon, D., Gonzalez, H., Updegraff, B.L., Orth, K. Vibrio parahaemolyticus type VI secretion system 1 is activated in marine conditions to target bacteria, and is differentially regulated from system 2. *PLoS One*. **8**(4), e61086 (2013).
- Sana, T.G., et al. Salmonella Typhimurium utilizes a T6SS-mediated antibacterial weapon to establish in the host gut. *Proceedings of the National Academy of Sciences of the United States of America*. **113**(34), E5044-E5051 (2016).
- Schwarz, S., et al., Burkholderia type VI secretion systems have distinct roles in eukaryotic and bacterial cell interactions. *PLoS Pathogens*. **6**(8), e1001068 (2010).
- Wenren, L.M., Sullivan, N.L., Cardarelli, L., Septer, A.N., Gibbs, K.A., Two independent pathways for self-recognition in Proteus mirabilis are linked by type VI-dependent export. *MBio*. **4**(4), e00374-13 (2013).
- García-Bayona, L., Guo, M.S., Laub, M.T.J.E. Contact-dependent killing by Caulobacter crescentus via cell surface-associated, glycine zipper proteins. *Elife*. **6**, 24869 (2017).
- Stubbendieck, R.M., Straight, P.D. Multifaceted interfaces of bacterial competition. *Journal of bacteriology*. **198**(16), 2145-2155 (2016).
- Cornforth, D.M., Foster, K.R. Antibiotics and the art of bacterial war. *Proceedings of the National Academy of Sciences of the United States of America*. **112**(35), 10827-10828 (2015).
- Shank, E.A., Kolter, R. New developments in microbial interspecies signaling. *Current Opinion in Microbiology*. **12**(2), 205-214 (2009).
- Roelofs, K.G., Coyne, M.J., Gentyala, R.R., Chatzidaki-Livanis, M., Comstock, L.E. Bacteroidales secreted antimicrobial proteins target surface molecules necessary for gut colonization and mediate competition in vivo. *MBio*. **7**(4), e01055-16 (2016).

14. Dey, A., Vassallo, C.N., Conklin, A.C., Pathak, D.T., Troselj, V., Wall, D. Sibling rivalry in *Myxococcus xanthus* is mediated by kin recognition and a polyploid prophage. *Journal of bacteriology*. **198**(6), 00964-15 (2016).
15. Danka, E.S., Garcia, E.C., Cotter, P.A. Are CDI systems multicolored, facultative, helping greenbeards? *Trends in Microbiology*. **25**(5), 391-401 (2017).
16. Willett, J.L., Ruhe, Z.C., Coulding, C.W., Low, D.A., Hayes, C.S. Contact-dependent growth inhibition (CDI) and CdiB/CdiA two-partner secretion proteins. *Journal of molecular biology*. **427**(23), 3754-3765 (2015).
17. Cianfanelli, F.R., Monlezun, L., Coulthurst, S.J. Aim, load, fire: the type VI secretion system, a bacterial nanoweapon. *Trends in Microbiology*. **24**(1), 51-62 (2016).
18. Joshi, A., Kostiuik, B., Rogers, A., Teschler, J., Pukatzki, S., Yildiz, F.H., Rules of engagement: the type VI secretion system in *Vibrio cholerae*. *Trends in microbiology*. **25**(4), 267-279 (2017).
19. Speare, L., et al., Bacterial symbionts use a type VI secretion system to eliminate competitors in their natural host. *Proceedings of the National Academy of Sciences of the United States of America*. **115**(36), E8528-E8537 (2018).
20. Shank, E.A., Using coculture to detect chemically mediated interspecies interactions. *Journal of Visualized Experiments*. **80**, (2013).
21. Long, R.A., Rowley, D.C., Zamora, E., Liu, J., Bartlett, D.H., Azam, F. Antagonistic interactions among marine bacteria impede the proliferation of *Vibrio cholerae*. *Applied and Environmental Microbiology*. **71**(12), 8531-8536 (2005).
22. Dunn, A.K., Millikan, D.S., Adin, D.M., Bose, J.L., Stabb, E.V. New rfp-and pES213-derived tools for analyzing symbiotic *Vibrio fischeri* reveal patterns of infection and lux expression in situ. *Applied and Environmental Microbiology*. **72**(1), 802-810 (2006).
23. Sana, T.G., et al., The second type VI secretion system of *Pseudomonas aeruginosa* strain PAO1 is regulated by quorum sensing and Fur and modulates internalization in epithelial cells. *Journal of Biological Chemistry*. **287**(32), 27095-27105 (2012).
24. Bachmann, V., Kostiuik, B., Unterweger, D., Diaz-Satizabal, L., Ogg, S., Pukatzki, S. Bile salts modulate the mucin-activated type VI secretion system of pandemic *Vibrio cholerae*. *PLoS*. **9**(8), e0004031 (2015).
25. Ishikawa, T., Rompikuntal, P.K., Lindmark, B., Milton, D.L., Wai, S.N. Quorum sensing regulation of the two hcp alleles in *Vibrio cholerae* O1 strains. *PloS One*. **4**(8), e6734 (2009).
26. Ishikawa, T., et al., Pathoadaptive conditional regulation of the type VI secretion system in *Vibrio cholerae* O1 strains. *Infection and immunity*. **80**(2), 575-584 (2012).
27. Pollack-Berti, A., Wollenberg, M.S., Ruby, E.G. Natural transformation of *Vibrio fischeri* requires tfoX and tfoY. *Environmental Microbiology*. **12**(8), 2302-2311 (2010).
28. Meibom, K.L., Blockesh, M., Dolganov, N.A., Wu, C.Y., Schoolnik, G.K. Chitin induces natural competence in *Vibrio cholerae*. *Science*. **310**(5755), 1824-1827 (2005).
29. Borgeaud, S., Metzger, L.C., Scrignari, T., Blockesh, M., The type VI secretion system of *Vibrio cholerae* fosters horizontal gene transfer. *Science*. **347**(6217), 63-67 (2015).
30. Townsley, L., Mangus, M.P.S., Mehic, S., Yildiz, F.H. Response of *Vibrio cholerae* to low-temperature shift: CpsV regulates type VI secretion, biofilm formation, and association with zooplankton. *Applied and Environmental Microbiology*. **82**(14), 00807-16 (2016).
31. Huang, Y., et al., Functional characterization and conditional regulation of the type VI secretion system in *Vibrio fluvialis*. *Frontiers in microbiology*. **8**, 528 (2017).